



POISON
KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS
BEFORE OPENING
OR USING

Thiss-Off®

THISTLE & BROADLEAF
HERBICIDE

**ACTIVE CONSTITUENTS: 750 g/L MCPA
(PRESENT AS THE DIMETHYLAMINE SALT)**

GROUP 4 HERBICIDE

For the selective control of broadleaf weeds in cereals, linseed, pastures,
rice, sugar cane and turf as per Directions for Use.

IMPORTANT: READ THE ATTACHED LEAFLET BEFORE USING THIS PRODUCT

WEED FORCE THISS-OFF THISTLE & BROADLEAF HERBICIDE

DIRECTIONS FOR USE

INDEX

RESTRAINTS.....	4
TABLE 1: FIELD CROPS	4
TABLE 2: PASTURES, SEED CROPS & TURF	8
TABLE 3: SPOT SPRAYING	12
TABLE 4: NON-CROP AREAS, FALLOW LAND, INDUSTRIAL & COMMERCIAL AREAS, RIGHTS-OF-WAY	14
WITHHOLDING PERIOD	14
WEEDS TABLE	15
GENERAL INSTRUCTIONS	20
APPLICATION/MIXING INFORMATION	20
RESISTANT WEEDS WARNING	21
EQUIPMENT MAINTENANCE AND USAGE	22
COMPATIBILITY/TANK MIXES	22
TANK MIXTURES	22
PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS	22
LEGUME TOLERANCE	22
PROTECTION OF LIVESTOCK	22
PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT	22
RE-ENTRY PERIOD	22
STORAGE AND DISPOSAL	22
SAFETY DIRECTIONS	23
FIRST AID	23
SAFETY DATA SHEET (SDS)	23
CONDITIONS OF SALE	23

WEEDFORCE Pty Ltd ACN 602 207 152
2072 Logan Rd Mt Gravatt QLD 4122
P: 0488 997 923 www.weedforce.com.au

APVMA Approval No:
83914/109240

© Registered Trademark of Weed Force Pty Ltd
© Weed Force Pty Ltd 2017

DIRECTIONS FOR USE

RESTRAINTS

DO NOT apply if crop or weeds are stressed due to dry or excessively moist conditions.

DO NOT apply if rain is likely within 6 hours.

1. FIELD CROPS

Situation & Crop	Weeds Controlled	State	Rate Volume/ha	Critical Comments	
Wheat, Barley, Oats, Triticale	Refer to Weed Table	SA, Tas only	460mL-1.45L	Apply through the tillering stage after the crop plants have 5 leaves and before they commence jointing.	
		Qld only		Apply from early tillering (4 to 5 fully emerged main stem leaves plus one or more tillers) until before the boot stage (visible swelling of the head at the top of the main stem).	
		NSW, ACT only		Apply 460mL rate after crop has reached 5 leaf stage and weeds marked E in Weed Table have 3-6 leaves. Higher rate after the first node can be felt at the base of a tiller and before swelling of the head can be felt in a tiller.	
		Vic only		460mL - 960mL	Apply 460mL rate after crop has reached 2-3 leaf stage and weeks marked E in Weed Table have 2-4 leaves. Higher rate after the crop plants have 5 leaves, to fully tillered.
		WA only		600mL - 1.35L	Apply to crop from 5 leaf to Flag leaf emergence. (Zadoks 15-37)
Wheat, Barley, Triticale, Cereal Rye	Capeweed, Double Gee, Erodium, Mustard, Rapistrum, Wild Turnip, Wild Radish	WA only	270mL plus 350mL diuron (500g/L)	CROP: 3-4 leaf stage (Zadoks 13-14)	
			330mL plus 400mL diuron (500g/L)	CROP: 4-5 leaf stage (Zadoks 14-15)	
	Amsinckia (Yellow Burr) Capeweed, Charlock, Deadnettle, Double Gee, (Spiny Emex, Three Corner Jack), Erodium spp, Mustard, Turnip Weed, White Ironweed (Corn Gromwell, Sheep Weed), Wild Radish, Wild Turnip	NSW, ACT, Vic only	330mL plus 500mL diuron (500g/L)	Apply by boom with 50-100L/ha water, when most weeds have germinated and are in 2-5 leaf stage. Crop and weeds should be dry at time of application. Some temporary yellowing of crop may occur after application. DO NOT apply to undersown medic or sub-clovers.	
	Capeweed, Mustard, Three Corner Jack, Turnip Weed, Wild Radish, Wild Turnip, Wards Weed, Common Ice Plant, Pimpernel	SA only	230mL plus 500mL diuron (500g/L)	DO NOT use on very light sandy soils.	

Situation & Crop	Weeds Controlled	State
Wheat, Barley Oats, Triticale	Soldier Thistle	SA only
	Skeleton Weed	

Rate Volume/ha	Critical Comments
460mL plus 200mL clopyralid	Apply to crop at tillering stage after the crop has 5 eaves and before the crop commences jointing.
660mL plus 500mL clopyralid	Apply to rosettes from 5-15cm diameter when the crop is between the 5 leaf and late tillering growth stages. It is essential that all the Skeleton Weed is emerged and minimum of 5cm diameter, which is usually not before August. This rate will provide control of Skeleton Weed until harvest and will substantially reduce seed numbers the following season.

Situation & Crop	Weeds Controlled	State	Weed Stage
Field Peas	Prickly Lettuce (<i>Lactuca serriola</i>)	NSW, Vic, Tas, SA only	4 leaf
			6 leaf
	Charlock (Wild Mustard) (<i>Sinapis arvensis</i>)		4 leaf
			6 leaf

Rate Volume/ha	Critical Comments
80mL plus 125mL of diflufenican (500g/L)	<p>FIELD PEAS: Apply early post-emergence after the third node stage and before the start of flowering.</p> <p>Warning: Field peas grown on high pH soils in the presence of free lime may be less tolerant to diflufenican (500g/L).</p> <p>APPLICATION AND WEED CONTROL Apply when weeds are actively growing. For optimum results apply 4 to 6 weeks post-sowing. Application beyond 8 weeks post-sowing may result in reduced levels of weed control. In most situations the rate specified for each weed size will give satisfactory control. Under certain conditions such as:</p> <ul style="list-style-type: none"> • high crop and weed density, • late season germinations, • abnormal weed growth (including early flowering); higher rates of product (up to the maximum rate of application specified for that weed) may be required. <p>Diflufenican (500g/L) will not effectively control;</p> <ul style="list-style-type: none"> • regrowth of suppressed weeds, • transplanted weeds, • regrowth from rhizomes or roots, • weeds growing under stress from previous herbicide applications.
100mL plus 150mL of diflufenican (500g/L)	
80mL plus 125mL of diflufenican (500g/L)	
100mL plus 150mL of diflufenican (500g/L)	

2. PASTURES, SEED CROPS & TURF

Situation & Crop	Weeds Controlled	State	Rate Volume/ha	Critical Comments
Linseed	Refer to Weed Table	Qld only	460mL - 1.45L	Apply when crop is 10-15cm tall with at least 170L/ha of water. DO NOT spray after buds appear. Some wilting effect may be seen following application but crop usually recovers quickly. DO NOT apply by aircraft.
		NSW, ACT, Vic, SA, Tas only	460mL – 730mL	
Sugar Cane	Blue Top, Chinese Burr, Flannel Weed, Gambia Pea, Bell Vine, Streaked Rattle Pod, Bindweed, Pink Convolvulus, Cupids Flower, Merremia Vine, Morning Glory	Qld only	930mL	Post Emergent: Apply as a directed inter-row spray. DO NOT exceed rates with phenoxy sensitive varieties.
	As above plus Fat Hen, Noogoora Burr		1.45L	
Oil-Seed Poppies (<i>Papaver somniferum</i>)	Fat Hen (<i>Chenopodium album</i>) Spear Thistle (<i>Cirsium vulgare</i>)	Tas only	1.0L	Use only in consultation with the poppy contracting company. Apply at least one week after petal fall after poppy seed has been set. Apply by aircraft in 30-40L of water per ha. This treatment will control late germinating weeds if they are green and at bud or early flowering stage. Spraying at this time will twist weeds below the crop canopy. DO NOT apply in horticultural areas where there is a risk of spray or vapour drift over susceptible crops.

Situation & Crop	Weeds Controlled	State	Rate Volume/ha	Critical Comments
Rice	Dirty Dora, Starfruit, Sedges, Buddah (Butter) Pea	NSW only	485mL to 970mL	Apply by aircraft at early/mid tillering stage of rice but not later than panicle initiation. Use 485 mL/ha at 35 days after aerial sowing (approx. 2 litres per rice plant). Use 970 mL/ha at 45 days or more after cereal sowing (3 tillers or more per rice plant). The crop should be sprayed before flowering of the weeds. Water Management: Before herbicide application, lower water level in the bays to at least two thirds of the weed growth for direct contact with MCPA 750. Weeds covered by water will not be controlled. Raise water depth to normal levels 24 hours following spraying.
Flax	Refer to Weed Table	Vic, Tas only	325mL	Apply to crop when 8-20cm high.
Grass Pastures	Refer to Weed Table	All states	460mL – 2.7L	Apply to established grass-based pastures. Damage may occur to Legumes if present. See 'Legume Tolerance.' NOTE: Capeweed, Paterson's Curse and Variegated Thistle may be poisonous to stock after spraying.
	Soldier Thistle	SA only	460mL - 1.45L	Use low rate on seedlings, with no more than 2 true leaves. Use 100 litres of water for best results. If clover is present do not exceed 575mL.
	Seedling and established Horehound	NSW, ACT only	2L	Apply in Autumn when Horehound plants are at the seedling stage. If sub clover or medic plants are present they will be damaged by this application.
Grass Seed Crops	Refer to Weed Table	Qld, SA, Tas, Vic, WA only	460mL - 2.7L	Apply to established grass seed crops.
Sub Clover Seed Crops	Refer to Weed Table	Vic, SA only	460mL - 960mL	DO NOT exceed rate specified. Spray in Autumn & early Winter.

Situation & Crop	Weeds Controlled	State
Pastures - Spray/ Graze Technique	Caltrop, Capeweed, Charlock, Mustards, Paterson's Curse, Shepherd's Purse, Saffron, Slender, Spear or Variegated Thistle, Turnip Weed, Wild Radish, Wild Turnip	NSW, ACT, Vic only Tas only
	Refer to Weed Table	WA only
	Capeweed, Dandelion, Seedling Dock, Three Corner Jack, Geranium, Erodium, Mustard, Annual Thistles, Turnip Weed, Wild Radish, Wild Turnip	SA only
	Soldier Thistle	
Turf	Refer to Weed Table	All States

Rate Volume/ha	Critical Comments
230mL – 930mL	<p>Spray actively growing 6-8 weeks old weeds. Introduce stock 7-10 days after spraying, preferably sheep as cattle are less effective. Use stocking rate at least 5 times heavier than normal until weeds have been reduced but before survival of desirable pasture species is threatened. Lucerne and medics may be damaged and should be grazed short before spraying. Refer 'Legume Tolerance.'</p> <p>Precaution: An increased quantity of poisonous plants may be eaten by stock using Spray/Graze, eg. Caltrop, Capeweed, Paterson's Curse and Variegated Thistle and deaths could result from causes such as nitrate poisoning. With Paterson's Curse preferably graze stock soon destined for slaughter and avoid extended periods of grazing. Avoid grazing with young or breeding stock.</p> <p>DO NOT graze horses or pigs on Paterson's Curse.</p> <p>Observe Withholding Period of at least 7 days after spraying.</p>
460mL	
610mL - 1.25L	
330mL	
395mL	
330mL or 2.8mL per 100 sq. metres	Newly laid - Delay until well established or at least two months old. Apply in high volume to actively growing weeds. See 'Legume Tolerance'. Buffalo grass is more easily damaged than other grasses
930mL – 1.8L or 10-19mL per 100 sq. metres	Established - Apply in high volume to actively growing weeds. DO NOT mow for 2 days before application. Some transitory damage may occur to fine turf grasses and strawberry clover. See 'Legume Tolerance'. Buffalo grass is more easily damaged than other grasses.

3. SPOT SPRAYING

Situation	State
High volume spraying	All states
Knapsack application	

Mixing Rate / Critical Comments
Add 130mL/100L of water for easy to kill weeds such as Wild Turnip, Mustards and Wild Radish. Add 270mL/100L of water for more difficult to control weeds such as Docks, Hoary Cress and Bindweed.
Add 13mL/10L of water for easy to kill weeds such as Wild Turnip, Mustards and Wild Radish. Add 27mL/10L of water for more difficult to control weeds such as Docks, Hoary Cress and Bindweed.

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION

4. NON-CROP AREAS, FALLOW LAND, INDUSTRIAL & COMMERCIAL AREAS, RIGHTS-OF-WAY

Situation & Crop	Weeds Controlled	State
Non-Crop Areas, Fallow Land, Industrial and Commercial Areas, Rights of Way	Refer to Weed Table	All states

Rate Volume/ha	Critical Comments
Refer to Weed Table	Apply at rates shown in Weed Table when weeds only are present and no crop or pasture is involved.

**NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER CONTRARY TO THIS LABEL
UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION**

WITHHOLDING PERIOD:

**CEREALS, PASTURE - DO NOT GRAZE OR CUT FOR STOCK FOR 7 DAYS AFTER APPLICATION.
HARVEST WITHHOLDING PERIOD: NOT REQUIRED WHEN USED AS DIRECTED Oilseed Poppies:
DO NOT HARVEST FOR 5 WEEKS AFTER APPLICATION**

WEEDS TABLE

Where weeds are to be sprayed in a crop or pasture, use only the rate given for the particular crop or situation indicated under the Directions for Use table. In most cases this will give control, however some hard to kill weeds or those in advanced stages of growth may only be suppressed. The rates listed in the Weed Table below are for use where weeds only are present and no crop or pasture is involved. Use at these rates in a crop or pasture other than for spot spraying may cause damage. This product should only be used in those States where a rate or range of rates is indicated for the particular weeds listed.

Refer to WEEDS TABLE pages 16 - 21

WEEDS TABLE**APPLICATION RATE PER HECTARE**

#: refers to Critical Comments for Wheat, Barley, Oats, and Triticale

Application Rate per Hectare					Application Rate per Hectare		
Weeds Controlled	VIC	NSW, ACT	SA	TAS	QLD	WA	Critical Comments
Ball Mustard	-	-	1.15L	-	-	-	Apply at rosette stage
Bathurst Burr	1.35L	660mL-1.35L	1.35L	1.35L	1.35L	1.35L	Spray young seedlings only
Black Bindweed	-	970mL-1.35L	-	-	930mL-1.35L	-	Apply at young stage prior to flowering
Californian Burr	-	660mL-1.35L	-	1.45L	-	-	Spray young seedlings only
Cape Tulip	-	-	-	-	-	1.35L	Apply just prior to flowering. Gives mild suppression only.
Capeweed	-	1.45L	1.45L	1.45L	1.45L	-	Apply at seedling stage
Carrot Weed	1.35L	-	-	-	1.35L	-	Apply at seedling stage
Charlock	460mL	660mL	460mL	660mL 1.45L	460mL	-	Spray up to rosette stage
Cutleaf Mignonette	-	-	1.45L	-	-	-	Apply at young rosette stage
Dandelion	1.35L	1.35L	1.35L	1.35L	1.35L	1.35L	Apply at young rosette stage
Deadnettle	-	1.45L	-	-	1.45L	-	Moderately susceptible. Apply at seedling stage.
Docks	690mL	-	1.45L	1.45L	-	-	Apply at rosette stage and before flower heads emerge. Only the top growth of old established plants will be affected.
Fat Hen	1.35L	660mL-1.35L	-	1.35L	1.35L	-	Spray up to rosette stage.
Fennel	1.8L	1.8L	1.8L	1.8L	1.8L	-	Apply at young stage
Field Bindweed	1.35L	-	1.35L	1.35L	1.35L	1.35L	Apply at bud stage, repeating over several seasons.
Field Cress	-	-	-	1.45L	-	-	Apply at young stage
Flatweed or Catsear	-	-	-	1.45L	-	930mL	Apply when weeds are 10-15cm high
Fumitory (red)	930mL	930mL	930mL	-	930mL	-	Apply at young stage prior to flowering
Hoary Cress	3L	-	1.45L	-	3L	-	Apply before flowering

WEEDS TABLE (CONTINUED)

Application Rate per Hectare				
Horehound	-	2L	1.45L	-
Hedge Mustard	730mL #	-	460mL	460mL
Hexham Scent or Melilotus	960mL	800mL	930mL	-
Lesser Swinecress	-	-	-	1.45L
Lincoln Weed	-	-	460mL	-
London Rocket	-	-	-	-
Lupins	-	460mL-960mL	-	-
Mintweed	1.35L	1.35L	-	-
Mustards	730mL #	660mL #	730mL	730mL
Noogoora Burr	1.35L	660mL-1.35L	1.35L	-
Opium Poppy	-	-	-	1.45L
Paterson's Curse	-	660-960mL	-	-
Pimpernel	-	-	460mL	-
Plantains	-	1.35L	-	1.35L
Rapeseed	-	660-960mL	-	-
Rough Poppy	-	460mL-960mL	-	-
Safflower	-	460-660mL	-	-
Shepherds Purse	-	-	-	1.45L
Skeleton Weed	-	960mL-1.35L	-	-
Sorrel	-	-	1.45L	-
Stinkwort	1.35L	660-960mL	1.35L	-
Sunflower (seedlings)	-	690mL-1L	-	-

Application Rate per Hectare		
-	-	Apply at rosette stage
460mL	825mL	Apply at rosette stage
930mL	-	Apply at young seedling stage
-	-	Apply at young stage
-	-	Apply at rosette stage
-	825mL	
-	825mL	Spray up to 10cm high
1.35L	-	Spray young seedlings only
-	825mL	Spray up to rosette stage
1.35L	1.35L	Spray young seedlings only
-	-	Apply at young stage
2.7L	-	Apply early rosette stage
-	-	Apply at rosette stage
1.35L	-	Apply at young stage
-	-	Spray up to rosette stage
-	-	Spray up to rosette stage
-	-	Apply at rosette stage
-	-	Spray rosettes before aerial growth commences
-	-	Apply to rosette stage
-	1.35L	Apply when weeds are 10-15cm high
-	-	Spray up to rosette stage

WEEDS TABLE (CONTINUED)

Application Rate per Hectare				
Thistles - Artichoke	1.05L	-	1.45L	-
- Nodding	-	1.65L	-	-
- Saffron	960mL-1.65L	660mL-1.35L	930mL-1.65L	930mL-1.65L
- Scotch or Cotton	1.45L		460-930mL	1.45L
- Slender	2L	660mL-1.35L	660mL	2L
- Soldier	730mL	-	730mL	-
- Spear	-	960mL-1.35L	-	1.45L
- Star	-	-	730-930mL	-
- Variegated	960mL	660mL-1.35L	460-930mL	960mL
Turnip Weed	730mL #	660mL #	660mL	-
Wild Artichoke	-		1.45L	-
Wild Radish	930mL	660mL #	930mL	1.35L
Wild Turnip	730mL #	660mL #	460mL	1.35L

GENERAL INSTRUCTIONS

Weeds should be sprayed while actively growing and at their most susceptible stage. As a general guide, annuals should be sprayed when young and perennials just prior to appearance of bud stage. Extremes of cold or drought are unfavourable. The best conditions are when the soil is moist, weather fine and rain unlikely within 6 hours of application. DO NOT spray crops or weeds outside the stages indicated in 'Critical Comments' as damage, loss of yield or inadequate weed control may result. DO NOT spray in high winds.

APPLICATION/MIXING INFORMATION

This product mixes readily with water. Half fill the spray tank with clean water and add the required amount of product. Agitate thoroughly before adding the remaining water.

- BOOM SPRAYING**
- Use 30-120 litres water per hectare.
 - Use minimum of 170 litres water per hectare for LINSEED.

Application Rate per Hectare		
1.35L	460mL	Apply at rosette stage
-	-	Spray early rosettes. Repeat spray required in following Spring or Autumn
930mL-1.65L	930mL-1.65L	Spray up to rosette stage (September WA)
		Apply at seedling stage
-	2L	Spray young seedlings only
-	-	Spray young seedlings. Use higher rate for larger weeds
-	-	Apply at rosette stage
-	-	Apply at seedling stage. Use higher rate for larger weeds
960mL	960mL	Spray young seedlings only. Use 1.35L/ha (SA) for well-developed weeds.
-	-	Spray up to rosette stage
-	-	Apply at rosette stage
460-730mL	800mL	Spray up to rosette stage
460-730mL	600mL	Spray up to rosette stage

AERIAL SPRAYING - Use 10-90 litres water per hectare.

RESISTANT WEEDS WARNING

GROUP	4	HERBICIDE
--------------	----------	------------------

Weed Force Thiss-Off Thistle & Broadleaf Herbicide is a member of the Phenoxy group of herbicides. Weed Force Thiss-Off Thistle & Broadleaf Herbicide as the Disruptors of plant cell growth mode of action. For weed resistance management Weed Force Thiss-Off Thistle & Broadleaf Herbicide is a Group 4 herbicide. Some naturally occurring weed biotypes resistant to Weed Force Thiss-Off Thistle & Broadleaf Herbicide and other disruptors of plant cell growth herbicides may exist through normal genetic variability in any weed population. The resistant individuals can eventually dominate the weed population if these herbicides are used repeatedly. These resistant weeds will not be controlled by Weed Force Thiss-Off Thistle & Broadleaf Herbicide or other disruptors of plant cell growth herbicides. Since the occurrence of resistant weeds is difficult to detect prior to use, Weed Force Pty Ltd accepts no liability for any losses that may result from the failure of Weed Force Thiss-Off Thistle & Broadleaf Herbicide to control resistant weeds.

EQUIPMENT MAINTENANCE AND USAGE

Equipment that has been used for this chemical should not be used for the application of other materials to sensitive plants unless it has been cleaned with suitable tank cleaner or well washed out with a 2% solution of cloudy ammonia followed by several clear water rinses.

COMPATIBILITY/TANK MIXES

This product can be tank mixed with diflufenican, metsulfuron-methyl, dicamba, bromoxynil (200 g/L), Flowable Diuron, clopyralid, Sertin, Sprayseed, Flowable Terbutryn, Tordon and Tribunil. The effectiveness of Imidan, Le Mat (omethoate 290 g/L) and Dimethoate may be reduced by alkaline hydrolysis if tank mixed with this product. For these situations the addition of LI-700 is recommended. When tank mixing add the required amount of compatible product (and surfactant if recommended) when the spray tank is nearly full and while continuously agitating. If left overnight, agitate thoroughly before spraying.

TANK MIXTURES

Read and follow all label directions including restraints, spray drift restraints, mandatory no-spray zones, critical comments, withholding periods, regional use restrictions and safety directions for the tank mix products.

PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS

DO NOT spray in high winds.

Drift Warning: DO NOT apply under weather conditions or from spraying equipment that may cause spray to drift onto nearby susceptible plants/crops, cropping lands or pastures. Avoid spray drift onto susceptible crops such as cotton, vines, tomatoes, vegetables, fruit trees, legume crops and pastures, oilseed crops or susceptible trees (eg. kurrajongs, belahs, eucalypts).

LEGUME TOLERANCE

The use of MCPA is preferred to 2,4-D Amine or Ester in situations where legumes are undersown in the crop.

Note: The tolerance of pasture legumes to applied product can vary with species, variety, growing conditions, stage of growth and companion crop cover. DO NOT spray cereal crops undersown with lucerne as damage may result.

PROTECTION OF LIVESTOCK

Low hazard to bees. May be applied at any time as recommended in the Directions for Use.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

DO NOT contaminate dams, rivers or streams with the product or used containers.

RE-ENTRY PERIOD

Hand weeding should not be performed for 7 days after application unless workers wear cotton overalls buttoned to the neck and wrist (or equivalent clothing), chemical resistant gloves and footwear.

STORAGE AND DISPOSAL

Store in the closed, original container in a dry, cool, well-ventilated area.
DO NOT store for prolonged periods in direct sunlight.

Triple-rinse containers before disposal. Add rinsings to spray tank. Do not dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush, or puncture and deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose, clear of waterways, desirable vegetation and tree roots, in compliance with relevant local, state or territory government regulations. Do not burn empty containers or product. For refillable containers: Empty contents fully into application equipment. Close all valves and return to point of supply for refill or storage.

SAFETY DIRECTIONS

Harmful if swallowed. Attacks eyes. Will irritate the skin. When opening the container, preparing spray and using the prepared spray wear cotton overalls buttoned to the neck and wrist and a washable hat, elbow-length PVC gloves and goggles. If product on skin, immediately wash area with soap and water. If product or spray in eyes, wash it out immediately with water. After use and before eating, drinking or smoking, wash hands, arms, and face thoroughly with soap and water. After each day's use wash gloves, goggles and contaminated clothing.

FIRST AID

If poisoning occurs, contact a Doctor or Poisons Information Centre. Phone Australia 13 11 26.

SAFETY DATA SHEET

The SDS contains important information relating to health and safety. This product should be used in conjunction with information on the SDS and users are advised to read the SDS before using this product. The most up to date SDS can be obtained from www.weedforce.com.au

CONDITIONS OF SALE

To the extent permitted by law, all conditions and warranties and statutory or other rights or action which buyer or any other user may have against Weed Force Pty Ltd ("Weed Force") or Seller are hereby excluded. Weed Force hereby gives Notice to Buyer and other users that it will not accept responsibility for any direct, indirect or consequential loss however arising (including negligence) from the sale, supply, use or application of this product or reliance on product information or advice provided by Weed Force or on its behalf. No warranty expressed or implied is given by Weed Force regarding its suitability or efficiency for any purpose for which it is used by the buyer, whether in accordance with the directions or not. Weed Force's liability shall, in all circumstances, be limited to replacement of the product or a refund of the purchase price paid therefor.



WEEDFORCE Pty Ltd ACN 602 207 152 
2072 Logan Rd Mt Gravatt QLD 4122
P: 0488 997 923 www.weedforce.com.au

® Registered Trademark of Weed Force Pty Ltd
© Weed Force Pty Ltd 2017

APVMA Approval No:
83914/109240